

CONTENTS

■ Bouaziz A., Kerzabi F., Brahmi B. Impact of the natural mortality (<i>M</i>) variability on the evaluation of the exploitable stock of sardine, <i>Sardina pilchardus</i> (Actinopterygii: Clupeiformes: Clupeidae) of the central region of the Algerian coast.	87
■ Tiralongo F., Baldacconi R. A conspicuous population of the long-snouted seahorse, <i>Hippocampus guttulatus</i> (Actinopterygii: Syngnathiformes: Syngnathidae), in a highly polluted Mediterranean coastal lagoon	99
■ Ayyildiz H., Emre Y., Ozen O., Yağcı A. Age and growth of <i>Capoeta erhani</i> (Actinopterygii: Cypriniformes: Cyprinidae) from the Menzelet Reservoir, Turkey	105
■ Kovalev K.V., Balashov D.A., Cherniak A.L., Lebedeva E.B., Vasil'eva E.D., Vasil'ev V.P. The karyotype of the Amu Darya sturgeon, <i>Pseudoscaphirhynchus kaufmanni</i> (Actinopterygii: Acipenseriformes: Acipenseridae)	111
■ Vo A.-T.E., Ashley M.C., Dikou A., Newman S.P. Fishery exploitation and stock assessment of the endangered Nassau grouper, <i>Epinephelus striatus</i> (Actinopterygii: Perciformes: Serranidae), in the Turks and Caicos Islands	117
■ Cengiz Ö., Ismen A., Ozekinci U. Reproductive biology of the spotted flounder, <i>Citharus linguatula</i> (Actinopterygii: Pleuronectiformes: Citharidae), from Saros Bay (northern Aegean Sea, Turkey)	123
■ Domagała J., Kirczuk L., Dziewulska K., Pilecka-Rapacz M. Annual development of gonads of pumpkinseed, <i>Lepomis gibbosus</i> (Actinopterygii: Perciformes: Centrarchidae) from a heated-water discharge canal of a power plant in the lower stretch of the Oder River, Poland	131
■ Kielpinski M., Kempter J., Panicz R., Keszka S. Genetic identifiability of selected populations of Indian mackerel, <i>Rastrelliger kanagurta</i> (Actinopterygii: Perciformes: Scombridae)—CELFIISH Project—Part 1	145
■ Piazzini S., Segos I., Favilli L., Manganelli G. The first European record of the Indonesian snakehead, <i>Channa micropeltes</i> (Actinopterygii: Perciformes: Channidae)	153
■ Mousavi-Sabet H., Khataminejad S., Vatandoust S. Length–weight and length–length relations of the seven endemic <i>Alburnus species</i> (Actinopterygii: Cypriniformes: Cyprinidae) in Iran	157
■ Erguden D., Turan C., Gurlek M., Uyan A., Reyhaniye A.N. First record of marbled stingray, <i>Dasyatis marmorata</i> (Elasmobranchii: Myliobatiformes: Dasyatidae), on the coast of Turkey, north-eastern Mediterranean	159
■ Oikonomou A., Anastasiadou Ch., Taskoudis Th., Leonardos I.D. Length–weight relations of seven native fish species (Actinopterygii) from the Louros River, Greece	163
■ Seyhaneyildiz Can S., Kutluyer F., Can E., Kayış Ş., Delihasan Sonay F., Köse Ö., Aksu Ö., Erdamar H., Yiğitoğlu M.R., Kayim M. Effect of dietary kefir on the digestive and liver enzymes activities, and glucose level of Coruh trout, <i>Salmo coruhensis</i> (Actinopterygii: Salmoniformes: Salmonidae)	167
■ EDITORIAL POLICY AND INSTRUCTIONS FOR AUTHORS	171

